

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 5/04, 15/00, 15/05, 15/29, 15/64, 15/82, A01H 3/00, 4/00, 5/00</b>		A1	(11) International Publication Number: <b>WO 00/26345</b>
			(43) International Publication Date: 11 May 2000 (11.05.00)
(21) International Application Number: <b>PCT/US99/25667</b>		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 3 November 1999 (03.11.99)			
(30) Priority Data: 09/185,244 3 November 1998 (03.11.98) US			
(71) Applicant (for all designated States except US): PLANT GENETIC SYSTEMS NV [BE/BE]; Jozef Plateastraat 22, B-9000 Ghent (BE).			
(72) Inventors; and (75) Inventors/Applicants (for US only): MICHIELS, Frank [BE/BE]; Poelstraat 291, B-9820 Bottelare (BE). JOHN-SON, Kirk [US/US]; 3926 Yana Place, Davis, CA 95616 (US).			
(74) Agents: LAWRENCE, William, F. et al.; Frommer Lawrence & Haug LLP, 745 Fifth Avenue, New York, NY 10151 (US).		<b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: <b>GLUFOSINATE TOLERANT RICE</b>			
(57) Abstract <p>This invention pertains to rice plants, plant material and seeds characterized by harboring a specific transformation event particularly by the presence of the <i>bar</i> gene under control of a CaMV 35S promoter, at a specific location in the rice genome. The rice plants of the invention combine glufosinate tolerance with optimal overall agronomic performance, genetic stability and adaptability to different genetic backgrounds.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Title of the invention**Glufosinate Tolerant Rice**5 Field of the invention

This invention pertains to rice plants, plant material and seeds characterized by harboring a specific transformation event particularly by the presence of the *bar* gene under control of a CaMV 35S promoter, at a specific location in the rice genome. The  
10 rice plants of the invention combine glufosinate tolerance with optimal overall agronomic performance, genetic stability and adaptability to different genetic backgrounds.

All documents cited herein are hereby incorporated herein by reference.

15

Background of the invention

The phenotypic expression of a transgene in a plant is determined both by the structure of the gene itself and by its location in the plant genome. At the same time  
20 the presence of the transgene at different locations in the genome will influence the overall phenotype of the plant. The agronomically or industrially successful introduction of a commercially interesting trait in a plant by genetic manipulation can be a lengthy procedure dependent on different factors. The actual transformation and regeneration of genetically transformed plants are only the first in a series of selection  
25 steps which include extensive genetic characterization, breeding, and evaluation in field trials.

Rice production is commonly threatened by various weeds. Some of these can be highly competitive and in cases of severe infestation can result in yield loss of such  
30 magnitude that it makes the crop economically unattractive. For direct-seeded, mechanized rice cultivation typical of temperate production, both cultural practices

(e.g. crop rotation, irrigation management) and herbicides are necessary to control weeds (Hill et al. 1994).

The *bar* gene (Thompson et al, 1987, EMBO J. 6:2519-2523; Deblock et al. 1987, EMBO J. 6:2513-2518) is a gene encoding the enzyme phosphinothricin acetyl transferase (PAT), which, when expressed in a plant, confers resistance to the herbicidal compounds phosphinothricin (also called glufosinate) or bialaphos (see also for example US patents, 5,646,024 and 5,561,236) and salts and optical isomers thereof. Other genes encoding PAT have been described (see for example: Wohlleben et al., 1988, Gene 70:25-37; EP 275,957; US 5,276,268; US 5,637, 489; US 5,273,894).

The transformation of monocotyledonous plants by electroporation of intact tissue capable of forming compact embryogenic callus or compact embryogenic callus obtained from such tissue is described in US Patent 5,641,664. Herein, transformation of compact embryogenic callus of rice by electroporation of a *bar* gene and the regeneration of transgenic rice plants is disclosed.

Transgenic rice plants containing the *gus* gene with either the *bar* gene, or the *hyg* gene conferring resistance to hygromycin, obtained by the transformation of cells of immature rice embryos by bombardment with DNA-coated gold particles have been described (Christou et al, 1991: Biotechnology 9:957).

Transformation of rice with the *bar* gene by electroporation of aggregated suspension cells is described in US Patent 5,679,558.

However, the foregoing documents fail to teach or suggest the present invention.

Summary of the invention

The present invention relates to a transgenic, glufosinate tolerant rice plant, cell, tissue or seed, which is characterized by one or both of the following characteristics:

- 5 a) the genomic DNA of the plant, cell, tissue or seed is capable of yielding one or more, such as at least two, advantageously at least three, preferably at least four, for instance at least five, more preferably six of the restriction fragments or pairs of restriction fragments selected from the group of:
  - 10 i) one EcoRI fragment with a length between about 1159 and about 1700 bp, preferably of about 1327 bp;
  - ii) one pair of BamHI fragments wherein one has a length between about 805 and about 1093 bp, preferably of about 805 bp and the other has a length between about 1700 and about 2140 bp, preferably of about 2,0 kbp;
  - 15 iii) one pair of EcoRV fragments wherein one has a length between about 2838 and about 4507 bp, preferably of about 3,8 kbp and the other has a length of more than about 5077 bp, preferably of about 12 kbp;
  - iv) one HindIII fragment with a length between about 5077 and about 11497 bp, preferably of about 5,3 kbp;
  - v) one pair of NcoI fragments both with lengths between about 2838 and about 4507  
20 bp, preferably one of about 3,1 kbp and one of about 4,1 kbp;
  - vi) one NsiI fragment with a length between about 4749 and about 11497 bp, preferably of about 5,1 kbp;
- 25 wherein each of the restriction fragments is capable of hybridizing under standard stringency conditions, with the about 1327 bp fragment obtainable by EcoRI digestion of the plasmid having the nucleotide sequence of SEQ ID No 8; and/or,
- b) the genomic DNA of the plant, cell, tissue or seed can be used to amplify a DNA fragment of between 290 and 350 bp, preferably of about 313 bp, using a polymerase chain reaction with two primers having the nucleotide sequence of SEQ ID No 4 and  
30 SEQ ID No 5, respectively (or includes a DNA fragment of about 290 to about 350 bp, preferably of about 313 bp amplified using a polymerase chain reaction with two

primers having the nucleotide sequence of SEQ ID No 4 and SEQ ID No 5, respectively).

The present invention relates to a transgenic, glufosinate tolerant rice plant, cell, tissue or seed, which is characterized in that the genomic DNA of the plant, cell, tissue or seed is capable of yielding at least one, advantageously at least two or more, for instance at least three, preferably at least four, for instance at least five, more preferably six of the restriction fragments or pairs of restriction fragments selected from the group described above comprising the restriction fragments or pairs of restriction fragments described under i), ii), iii), iv), v) and vi) above, whereby the selection can include any combination of i), ii), iii), iv), v) and vi) described above.

The present invention relates to a transgenic, glufosinate tolerant rice plant, cell, tissue or seed which is preferably characterized by both of the characteristics described under a) and b) above.

15

The invention also relates to the seed deposited at the ATCC under number ATCC 203352, a plant which is grown from this seed, and cells or tissues from a plant grown from this seed. The invention further relates to plants obtainable by propagation of, and/or breeding with a rice plant grown from the seed deposited at the ATCC under number ATCC 203352.

20

The invention further relates to plants, seeds, cells or tissues (e.g., rice plants, seeds, cells or tissues) comprising herein discussed flanking regions with the 35S-bar gene (as herein discussed) therebetween, or plants, seeds, cells, or tissues (e.g., rice plants, seeds, cells or tissues) comprising a nucleotide sequence which is at least 65%, e.g., at least 75%, such as at least 80%, for instance at least 85%, such as at least 90%, for example at least 95% or even 97% or 100% similar to a sequence disclosed herein, such as the sequence for the flanking region-35S-bar gene-flanking region construct, or the insertion region.

25

The invention further relates to a process for cultivating rice plants of the invention as described above, more particularly a process which comprises applying a herbicide with glufosinate as an active ingredient to the cultivated rice plants.

- 5 It is believed that the rice plants of the invention, when cultivated according to the process described above, which comprises applying a herbicide with glufosinate as an active ingredient, display improved growth as compared to untransformed rice of the same cultivar (US 5,739,082). Thus, the invention can comprehend a method for improving the yield or growth of rice plants

10

The invention also provides a process for breeding rice which comprises a crossing with the rice plants of the invention.

- 15 The invention further provides a process for producing a transgenic cell of a rice plant or a plant obtained therefrom, which comprises inserting a recombinant DNA molecule into a part of the chromosomal DNA of a rice cell characterized by the sequence of SEQ ID No 9 and, optionally, regenerating a rice plant from the transformed rice cell.

- 20 The invention further relates to a method for identifying a transgenic plant, or cells or tissues thereof, which method comprises establishing one or both of the following characteristics of the genomic DNA of the transgenic plant, or its cells or tissues:

a) the genomic DNA of the plant, cell, tissue or seed is capable of yielding at least three, preferably at least four, for instance at least 5, more preferably six of the restriction fragments or pairs of restriction fragments selected from the group of:

25

i) one EcoRI fragment with a length between 1159 and 1700 bp, preferably of about 1327 bp;

ii) one pair of BamHI fragments wherein one has a length between 805 and 1093 bp, preferably of about 805 bp and the other has a length between

30

1700 and 2140 bp, preferably of about 2,0 kbp;

- iii) one pair of EcoRV fragments wherein one has a length between 2838 and 4507 bp, preferably of about 3,8 kbp and the other has a length of more than 5077 bp, preferably of about 12 kbp;
- iv) one HindIII fragment with a length between 5077 and 11497 bp, preferably of about 5,3 kbp;
- v) one pair of NcoI fragments both with lengths between 2838 and 4507 bp, preferably one of about 3,1 kbp and one of about 4,1 kbp;
- vi) one NsiI fragment with a length between 4749 and 11497 bp, preferably of about 5,1 kbp;
- wherein each of the restriction fragments is capable of hybridizing under standard stringency conditions, with the 1327 bp fragment obtainable by EcoRI digestion of the plasmid having the nucleotide sequence of SEQ ID No. 8; and/or,
- b) the genomic DNA of the plant, cell, tissue or seed can be used to amplify a DNA fragment of between 290 and 350 bp, preferably of about 313 bp, using a polymerase chain reaction with two primers having the nucleotide sequence of SEQ ID No. 4 and SEQ ID No. 5, respectively.

The invention further relates to a kit for identifying the transgenic plants comprising the elite event of the present invention, said kit comprising PCR probes recognizing the foreign DNA and the 3' or 5' flanking sequence of GAT-OS2, preferably having the nucleotide sequence of SEQ ID No. 4 and SEQ ID NO. 5, respectively, for use in the PCR identification protocol.

#### Brief description of the drawings

The following detailed description, given by way of example, but not intended to limit the invention to specific embodiments described, may be understood in conjunction with the accompanying Figures, incorporated herein by reference, in which:

#### **Fig. 1. Restriction map obtained after digestion of GAT-OS2 genomic DNA**

Loading sequence of the gel analyzed by Southern blot: lane 1, Lambda DNA digested with PstI, lane 2, GAT-OS2 DNA digested with EcoRI, lane 3, GAT-OS2



7

DNA digested with BamHI, lane 4, GAT-OS2 DNA digested with EcoRV, lane 5, GAT-OS2 DNA digested with HindIII, lane 6, GAT-OS2 DNA digested with NcoI, lane 7, GAT-OS2 DNA digested with NsiI, lane 8, non-transgenic rice DNA, lane 9, Control plasmid DNA digested with EcoRI.

5

**Fig. 2. PCR analysis of different lines using the GAT-OS2 PCR identification protocol.** Loading sequence of the gel: lane 1, molecular weight marker (100bp ladder), lanes 2 to 11, DNA samples from rice plants comprising different transgenic events, lane 12, DNA from M202 wild-type, lane 13, DNA from Bengal wild-type, lane 14, negative control (water), lane 15, molecular weight marker (100bp ladder).

10

#### Detailed description

The term "gene" as used herein refers to any DNA sequence comprising several operably linked DNA fragments such as a promoter and a 5' untranslated region (the 5'UTR), which together form the promoter region, a coding region (which may or may not code for a protein), and an untranslated 3' region (3'UTR) comprising a polyadenylation site. Typically in plant cells, the 5'UTR, the coding region and the 3'UTR are transcribed into a RNA which, in the case of a protein encoding gene, is translated into the protein. A gene may include additional DNA fragments such as, for example, introns. As used herein, a genetic locus is the position of a given gene in the genome of a plant.

15

20

The term "chimeric" when referring to a gene or DNA sequence is used to refer to the fact that the gene or DNA sequence comprises at least two functionally relevant DNA fragments (such as promoter, 5'UTR, coding region, 3'UTR, intron) that are not naturally associated with each other and originate, for example, from different sources. "Foreign" referring to a gene or a DNA sequence with respect to a plant species is used to indicate that the gene or DNA sequence is not naturally found in that plant species.

25

30

- As used herein the term "transgene" refers to a recombinant DNA molecule as incorporated in the genome of a plant. The term "recombinant DNA molecule" is used to exemplify and thus can include an isolated nucleic acid molecule which can be DNA and which can be obtained through recombinant or other procedures. This
- 5 recombinant DNA molecule usually comprises at least one copy of at least one "gene of interest" (e.g. a chimeric gene) which is capable of conferring one or more specific characteristics to the transformed plant. A "transgenic plant" refers to a plant comprising a transgene in the genome of all of its cells.
- 10 The incorporation of a recombinant DNA molecule in the plant genome typically results from transformation of a cell or tissue (or from another genetic manipulation). The particular site of incorporation is either due to chance or is at a predetermined location (if a process of targeted integration is used).
- 15 The transgene can be characterized by the location and the configuration at the site of incorporation of the recombinant DNA molecule in the plant genome. The site in the plant genome where a transgene has been inserted is also referred to as the "insertion site" or "target site". Insertion of the transgene into the plant genome can be associated with a deletion of plant DNA, referred to as "target site deletion". A
- 20 "flanking region" or "flanking sequence" as used herein refers to a sequence of at least 20 bp, preferably at least 50 bp, and up to 5000 bp of the plant genome which is located either immediately upstream of and contiguous with or immediately downstream of and contiguous with the transgene. Transformation procedures leading to random integration of the transgene will result in transformants with different
- 25 flanking regions, which are characteristic and unique for each transformant. When the transgene is introduced into a plant through traditional crossing, its insertion site in the plant genome, or its flanking regions will generally not be changed. An "insertion region" as used herein refers to the region corresponding to the region encompassed by the insertion site (and possible target site deletion), the upstream and the
- 30 downstream flanking regions of a transgene in the (untransformed) plant genome.

Expression of the transgene is used to indicate that the gene(s) of interest comprised in the transgene is expressed so as to confer on the plant one or more phenotypic traits (e.g. herbicide tolerance) that were intended to be conferred by the introduction of the recombinant DNA molecule – the transforming DNA - used during transformation  
5 (on the basis of the structure and function of part or all of the gene(s) of interest).

An event is defined as a (artificial) genetic locus that, as a result of genetic manipulation, carries a transgene comprising at least one copy of a gene of interest. The typical allelic states of an event are the presence or absence of the transgene. An  
10 event is characterized phenotypically by the expression of the transgene. At the genetic level, an event is part of the genetic makeup of a plant. At the molecular level, an event is characterized by the restriction map (e.g. as determined by Southern blotting) and/or by the upstream and/or downstream flanking sequences of the transgene, and/or the molecular configuration of the transgene. Usually  
15 transformation of a plant with a transforming DNA comprising at least one gene of interest leads to a multitude of events, each of which is unique.

An elite event, as used herein, is an event which is selected from a group of events obtained by transformation with the same transforming DNA, based on the expression  
20 and stability of the transgene and its compatibility with optimal agronomic characteristics of the plant comprising it. Thus the criteria for elite event selection are one or more, preferably two or more, advantageously all of the following:

- a) That the presence of the transgene does not compromise other desired characteristics of the plant, such as those relating to agronomic performance or  
25 commercial value;
- b) That the event is characterized by a well defined molecular configuration which is stably inherited and for which appropriate diagnostic tools for identity control can be developed;
- c) That the gene(s) of interest in the transgene shows a correct, appropriate and stable  
30 spatial and temporal phenotypic expression, both in heterozygous (or hemizygous) and homozygous condition of the event, at a commercially acceptable level in a range

of environmental conditions in which the plants carrying the event are likely to be exposed in normal agronomic use;

It is preferred that the transgene is associated with a position in the plant genome that  
5 allows introgression into desired commercial genetic backgrounds.

The status of an event as an elite event is confirmed by introgression of the elite event in different relevant genetic backgrounds and observing compliance with one, two or all of the criteria e.g. a), b) and c) above.

10 An "elite event" thus refers to a genetic locus comprising a transgene, which answers to the above-described criteria. A plant, plant material or progeny such as seeds can comprise the elite event in its genome.

The "diagnostic tools" developed to identify an elite event or the plant or plant  
15 material comprising an elite event, are based on the specific genomic characteristics of the elite event, such as, a specific restriction map of the genomic region comprising the transgene and/or the sequence of the flanking region(s) of the transgene. A "restriction map" as used herein refers to a set of Southern blot patterns obtained after cleaving plant genomic DNA with a particular restriction enzyme, or set of  
20 restriction enzymes and hybridization with a probe sharing sequence similarity with the transgene (under specific conditions). Due to the (endogenous) restriction sites present in a plant genome prior to incorporation of the transgene, insertion of a transgene will alter the specific restriction map of that genome. Thus, a particular transformant or progeny derived thereof can be identified by one or more specific  
25 restriction patterns. The conditions for determining the restriction map of an event are laid out in a restriction map identification protocol.

Alternatively, plants or plant material comprising an elite event can be identified by testing according to a PCR identification protocol. This is a PCR using primers which  
30 specifically recognize the elite event. Essentially, a set of primers is developed which recognizes a) a sequence within the 3' or 5' flanking sequence of the elite event and

b) a sequence within the foreign DNA, which primers amplify a fragment (integration fragment) preferably of between 100 and 350 nucleotides. Preferably, a control is included of a set of primers which amplifies a fragment within a housekeeping gene of the plant species (preferably a fragment which is larger than the amplified integration fragment). The optimal conditions for the PCR, including the sequence of the specific primers is specified in a PCR identification protocol.

The term "similarity", for instance, with respect to a nucleotide sequence, is intended to indicate a quantitative measure of homology between two sequences. The percent sequence similarity can be calculated as  $(N_{ref} - N_{dif}) * 100 / N_{ref}$ , wherein  $N_{dif}$  is the total number of non-identical residues in the two sequences when aligned and wherein  $N_{ref}$  is the number of residues in one of the sequences. Hence, the DNA sequence AGTCAGTC will have a sequence similarity of 75% with the sequence AATCAATC ( $N_{ref} = 8$ ;  $N_{dif} = 2$ ). The invention comprehends nucleic acid molecules and with sequences having at least 65%, e.g., at least 70%, such as at least 75%, or at least 80% or advantageously at least 85%, for instance at least 90%, such as at least 95% or even 97% or 100% similarity with sequences disclosed herein, as well as plants, cells, tissues, seeds, and progeny thereof (e.g., rice plants, cells, tissues, seeds and progeny thereof) comprising such nucleic acid molecules. Alternatively or additionally, "similarity" with respect to sequences refers to the number of positions with identical nucleotides divided by the number of nucleotides in the shorter of the two sequences wherein alignment of the two sequences can be determined in accordance with the Wilbur and Lipmann algorithm (Wilbur and Lipman, 1983 PNAS USA 80:726) using a window size of 20 nucleotides, a word length of 4 nucleotides, and a gap penalty of 4, and computer-assisted analysis and interpretation of the sequence data including alignment can be conveniently performed using programs of the Intelligenetics™ Suite (Intelligenetics Inc. CA). Sequences which are "essentially similar" have a sequence similarity or identity of at least about 75%, advantageously at least about 80%, such as at least about 85%, preferably at least about 90%, especially about 95%, such as at least 97%, and especially about 100%. It is clear that when RNA sequences

are said to be essentially similar or similar, or have a degree of sequence identity with DNA sequences, thymidine (T) in the DNA sequence is considered equal to uracil (U) in the RNA sequence.

5 The present invention relates to the development of an elite event in rice, GAT-OS2, and the plants, plant cells, or plant material derived from this event. Plants comprising elite event GAT-OS2 were obtained through transformation with a 1501 bp PvuI-HindIII fragment of plasmid pB5/35Sbar (SEQ ID No 8) as described in example 1. The recombinant DNA molecule used for generation of this elite event comprises a

10 DNA sequence encoding the enzyme phosphinothricin acetyl transferase and the 35S promoter of Cauliflower Mosaic Virus, wherein the sequence encoding phosphinothricin acetyl transferase is under the control of the 35S promoter of Cauliflower Mosaic Virus (termed the "35S-bar gene"). The 35S promoter has a "constitutive" expression pattern in rice (Battraw et al, 1990, Plant Mol Biol 15:527-

15 538), which means that it is significantly expressed in most plant cell types, during most of the plant life cycle. The expression of the 35S-bar gene in rice plants confers resistance to herbicidal compounds phosphinothricin or bialaphos or glufosinate or more generally, glutamine synthetase inhibitors, or salts or optical isomers thereof.

20 Plants or plant material comprising GAT-OS2 can be identified according to the restriction map identification protocol described in Example 3b)(1) herein. Briefly, rice genomic DNA is digested with a selection (preferably at least one such as at least two, advantageously at least three, or at least four, or at least five such as three to six) of the following restriction enzymes: EcoRI, BamHI, EcoRV, HindIII, NcoI, NsiI, is

25 then transferred to nylon membranes and hybridized with the about 1327 bp EcoRI fragment of plasmid pB5/35Sbar. It is then determined for each restriction enzyme used whether the following fragments can be identified:

- EcoRI: one fragment of between about 1159 and about 1700 bp, preferably of about

1327 bp

- BamHI: one fragment of between about 1700 and about 2140 bp, preferably of about 2,0 kbp and one fragment of between about 805 and about 1093 bp, preferably of about 805 bp
- EcoRV: one fragment of more than about 5077 bp, preferably of about 12 kbp and one fragment of between about 2838 and about 4507 bp, preferably of about 3,8 kbp
- HindIII: one fragment of between about 5077 and about 11497 bp, preferably of about 5,3 kbp
- NcoI: two fragments of between about 2838 and about 4507 bp, preferably one of about 4,1 kbp and one of about 3,1 kbp
- NsiI: one fragment of between about 4749 and about 11497 bp, preferably of about 5,1 kbp.

The lengths of the DNA fragments are determined by comparison with a set of DNA fragments of known length, particularly the PstI fragments of phage lambda DNA.

- 15 If the plant material after digestion with at least one such as at least two, advantageously at least three, preferably at least four, especially with at least 5, more preferably with all of these restriction enzymes, yields DNA fragments with the same length as those described above, the rice plant is determined to harbor elite event GAT-OS2.

20

- Plants or plant material comprising GAT-OS2 can also be identified according to the PCR identification protocol described in Example 3b)(2) herein. Briefly, rice genomic DNA is amplified by PCR using a primer which specifically recognizes a flanking sequence of GAT-OS2, particularly the primer with the sequence of SEQ ID No 5, and a primer which recognizes a sequence in the transgene, particularly the primer with the sequence of SEQ ID No 4. Endogenous rice primers are used as controls. If the plant material yields a fragment of between about 290 and about 350 bp, preferably of about 313 bp, the rice plant is determined to harbor elite event GAT-OS2.

30

Plants harboring GAT-OS2 are also characterized by their glufosinate tolerance, which in the context of the present invention includes that plants are tolerant to the herbicide Liberty<sup>TM</sup>. Tolerance to Liberty<sup>TM</sup> is defined by the criterium that spraying of the plants in the three to four leaf stage (3V to 4V) with at least 200 grams active ingredient/hectare (g.a.i./ha), preferably 400 g.a.i./ha, and possibly up to 1600 g.a.i./ha, does not kill the plants. Plants harboring GAT-OS2 can further be characterized by the presence in their cells of phosphinothricin acetyl transferase as determined by a PAT assay (De Block et al, 1987, supra).

Plants harboring GAT-OS2 can, for example, be obtained from seeds deposited at the ATCC under number ATCC 203352. Such plants can be further propagated and/or used in a conventional breeding scheme to produce more transformed plants with the same characteristics or to introduce the elite event of the invention into other cultivars of the same plant species. Seeds obtained from these plants contain the elite event stably incorporated into their genome.

The rice plants of this invention can be cultivated in a conventional way. The presence of the transgene ensures that they are tolerant to glufosinate. Therefore, weeds in the fields where such rice plants are grown can be controlled by application of herbicides comprising glufosinate as an active ingredient (such as Liberty<sup>TM</sup>).

Plants harboring GAT-OS2 are also characterized by having agronomical characteristics which are comparable to the following commercially available rice varieties in the US: Priscilla, Cypress, Bengal, Cocadrie, Jefferson, Madison, M202, M201, M103, Drew, Kaybonnet, Lagrue. The agronomical characteristics of relevance are: plant height, strength/stiffness of straw, resistance to lodging, leaf morphology (length, width, and angle for flag leaf), time to maturity, floret configuration, panicle fertility, complete closure of the hull on the seed, grain size and shape, and grain production and yield.



It has been observed that the presence of the transgene in this region of the rice plant genome, more particularly at this site in the rice plant genome, confers particularly interesting phenotypic and molecular characteristics to this event. More specifically, the presence of a transgene at this particular site in the genome results in stable phenotypic expression of the transgene without significantly compromising any aspect of desired agronomic performance of the plant. Thus, the insertion region, corresponding to SEQ ID No 9, more particularly the insertion site of GAT-OS2 therein, is shown to be particularly suited for the introduction of a gene(s) of interest, such as a herbicide resistance gene, more specifically a gene encoding phosphinothricin acetyl transferase under the control of a 35S promoter, particularly the PvuI-HindIII fragment of plasmid pB5/35Sbar.

A recombinant DNA molecule can be specifically inserted in this insertion region by targeted insertion methods. Such methods are well known to those skilled in the art and comprise, for example, homologous recombination using a recombinase such as, but not limited to either FLP recombinase from *Saccharomyces cerevisiae* (US Patent 5,527,695), the CRE recombinase from *Escherichia coli* phage P1 (published PCT application WO 9109957, the recombinase from pSRI of *Saccharomyces rouxii* (Araki et al. 1985, J Mol Biol 182:191-203), or the lambda phage recombination system such as described in US Patent 4,673,640.

DNA can be inserted into a plant genome, such as a rice genome by techniques including, electroporation methods, bombardment with DNA-coated gold particles or biolistic methods, or agrobacterium or polyethylene glycol mediated methods, and the like

As used herein "comprising" is to be interpreted as specifying the presence of the stated features, integers, steps or components as referred to, but does not preclude the presence or addition of one or more features, integers, steps or components, or groups thereof. Thus, e.g., a nucleic acid or protein comprising a sequence of nucleotides or amino acids, may comprise more nucleotides or amino acids than the actually cited

ones, i.e., be embedded in a larger nucleic acid or protein. A chimeric gene comprising a DNA sequence which is functionally or structurally defined, may comprise additional DNA sequences, etc.

- 5 The following examples describe the development and characteristics of rice plants harboring the elite event GAT-OS2.

Unless otherwise stated, all recombinant DNA techniques are carried out according to standard protocols as described in Sambrook et al. (1989) *Molecular Cloning: A*  
10 *Laboratory Manual*, Second Edition, Cold Spring Harbour Laboratory Press, NY and in Volumes 1 and 2 of Ausubel et al. (1994) *Current Protocols in Molecular Biology*, Current Protocols, USA. Standard materials and methods for plant molecular work are described in Plant Molecular Biology Labfax (1993) by R.D.D. Croy published by BIOS Scientific Publications Ltd (UK) and Blackwell Scientific Publications, UK.

15

In the description and examples, reference is made to the following sequences:

- |                 |   |
|-----------------|---|
| SEQ ID No 1:    | sequence comprising the 5' flanking region of GAT-OS2             |
| SEQ ID No 2:    | sequence comprising the 3' flanking region of GAT-OS2             |
| 20 SEQ ID No 3: | sequence comprising the insertion site of GAT-OS2                 |
| SEQ ID No 4:    | OSA03: primer of the PCR identification protocol                  |
| SEQ ID No 5:    | OSA04: GAT-OS2-specific primer of the PCR identification protocol |
| SEQ ID No 6:    | OSA01 rice endogenous primer                                      |
| 25 SEQ ID No 7: | OSA02 rice endogenous primer                                      |
| SEQ ID No 8:    | plasmid pB5/35Sbar  |
| SEQ ID No 9:    | insertion region  |

**EXAMPLES**Example 1. Transformation of rice with a gene encoding herbicide resistance

- 5 a) Construction of the chimeric DNA comprising the bar gene under the control of a 35S promoter (pB5/35Sbar)

A plasmid pB5/35Sbar was constructed following standard procedures. The sequence of plasmid pB5/35Sbar is given in SEQ ID No 8. Digestion with PvuI-HindIII yielded  
 10 a 1501 bp fragment which comprised the following genetic elements:

Nucleotide coordinates	Genetic elements
2140-2195	Sequence derived from pUC19 (Yanish-Perron et al., Gene 33:103-119, 1985)
2196-2204	Synthetic polylinker sequence
2205-2398	Complement of 35S terminator (T35S) from Cauliflower Mosaic Virus (Franck A. et al., Cell 21:285-294, 1980; Pietrzak M. et al., Nucl. Acids Res. 14:5857-5868, 1986)
2399-2417	Synthetic polylinker sequence
2418-2969	Complement of bar gene from <i>Streptomyces hygroscopicus</i> (Thompson et al., EMBO J. 6:2519-2523, 1986)
2970-2985	Synthetic polylinker sequence
2986-3517	Complement of 35S promoter (P35S) from Cauliflower Mosaic Virus (Franck et al. above; Pietrzak et al., above)
3518-3641	Sequence derived from pUC19, (Yanisch-Perron et al., above)

The 1501 bp PvuI-HindIII fragment was purified by extraction of this fragment after electrophoresis.

## b) Transformation of rice

The Bengal Variety is a medium-grain rice developed by the Rice Research Station of the Louisiana Agricultural Experiment Station. The variety was officially released in 1992. The pedigree includes MARS and M201 (Linscombe S.D. et al. 1993, Crop Science: 33:645-646).

Transformation of rice plants with the 1501 bp PvuI-HindIII fragment of pB5/35Sbar was performed using direct DNA transfer. Selection was done on phosphinothricin (PPT) at all stages except plantlet regeneration, which was done in the absence of PPT to accelerate growth. This resulted in a set of primary transformants (plants of generation T<sub>0</sub>).

Example 2. Development of events

## a) Development of transgenic homozygous lines

The various T<sub>0</sub> hemizygous plantlets were transitioned from tissue culture, transferred to greenhouse soil, and allowed to flower and set seed. Plantlets were evaluated for fertility, fecundity and tolerance to glufosinate ammonium. 19 plants were selected for further analysis. T<sub>1</sub> seed produced by selfing was collected from these plants and grown in the field. T<sub>1</sub> plants were sprayed with Liberty™ herbicide at 800 grams active ingredient per hectare (g.a.i./ha; recommended dosage for farmers is 400 g.a.i./ha). The events that survived the herbicide application and segregated 3:1 for herbicide tolerance were selected for further evaluation. Tolerant plants were evaluated for damage (leaf tip burn).

T<sub>2</sub> seeds were harvested from panicles of all tolerant plants of selected events. These were sown in rows and T<sub>2</sub> plants were sprayed with Liberty™ herbicide (1600 g.a.i./ha) to evaluate segregation of the herbicide tolerance. Those rows that had 100% survivors and thus corresponded to lines which were homozygous for the transgene

were selected. These were again evaluated for herbicide damage and phenotypic traits. Further selection of events was made based on uniformity of phenotype within the panicle row (for the desired characteristics).

5 b) Characterization of transgenic events – selection of GAT-OS2

Transgenic events were further characterized for southern blot patterns, general phenotype and agronomic performance, and yield. Where appropriate these characteristics were determined in field conditions.

10

**Southern blot analysis**

Presence of the transgene was checked by standard Southern blot analysis using enzymatic digestion of rice genomic DNA with EcoRV and hybridization to the 1327 bp EcoRI fragment of pB5/35Sbar. The relative band intensity provided an indication on whether plants were homozygous or hemizygous for the transgenic locus. Two events were found to have simple insertions. This was confirmed by the fact that the segregation pattern of the transgene could be explained by Mendelian inheritance of a simple locus.

20

**General plant phenotype and agronomic performance**

T<sub>1</sub> and T<sub>2</sub> plants were evaluated for a number of phenotypic traits including plant height, strength/stiffness of straw, tendency to lodge, leaf morphology (too thin or incorrect angle for flag leaf), late maturity, floret configuration, panicle sterility or incomplete fertility, incomplete closure of the hull on the seed (which would lead to increased disease susceptibility), grain size and shape, and grain production and yield. Lines were evaluated to be similar (or improved) in displayed agronomic characteristics compared to the untransformed Bengal cultivar and the following rice varieties: Priscilla, Cypress, Cocadrie, Jefferson, Madison, M202, M201, M103, Drew, Kaybonnet, Lagrue. In some cases, the plants within a panicle row segregated

30

for somaclonal variation for one or more of the above-mentioned traits. Unless this resulted in the introduction of a commercially interesting phenotypic trait, these plants were discarded.

## 5 Field trials for yield evaluation

T<sub>2</sub> seeds were harvested in bulk from the selected homozygous populations and were compared to variety standards of Bengal. The seeds were planted as panicle rows in isolated blocks representing each event. Transgenic plots were sprayed with 1,600  
10 g.a.i./ha of Liberty<sup>TM</sup> herbicide or not sprayed ("no-spray" plots). Plots with non-transgenic variety standards were not sprayed with Liberty<sup>TM</sup>. Standard herbicide treatments to control local weeds were applied to all plots.

Transgenic events were tested for yield performance in different locations including  
15 Louisiana and Puerto Rico (winter nursery).

Statistical analysis of the agronomic parameters and ranking statistics of the plant morphology and other non-parametric data were completed to identify the best commercial candidate to compete with the parent variety, Bengal and the following  
20 rice varieties: Priscilla, Cypress, Cocadrie, Jefferson, Madison, M202, M201, M103, Drew, Kaybonnet, Lagrue. GAT-OS2 was the event showing the most utility for producing a range of breeding lines.

### Example 3. Characterization of event GAT-OS2

25

#### a) In-depth molecular and genetic analysis of the locus

Once the GAT-OS2 event was identified as the event in which expression of the transgene as well as overall agronomic performance were optimal, the locus of the  
30 transgene was analyzed in detail on a molecular level. This included detailed Southern blot analysis and sequencing of the flanking regions of the transgene.

(1) Southern blot analysis using multiple restriction enzymes

5 Leaf tissue was harvested from transgenic and control plants. Total genomic DNA was isolated from leaf tissue according to Dellaporta et al. (1983, Plant Molecular Biology Reporter, 1, vol.3, p.19-21). The DNA concentration of each preparation was determined by measuring the optical density in a spectrophotometer at a wavelength of 260 nm.

10 10 µg of genomic DNA was digested with restriction enzyme in a final reaction volume of 40 µl, applying conditions proposed by the manufacturer. The time of digestion and/or amount of restriction enzyme were adjusted to ensure complete digestion of the genomic DNA samples without non-specific degradation. After digestion, 4 µl of loading dye was added to the digested DNA samples, and they were  
15 loaded on a 1% agarose gel.

The following control DNAs were also loaded on the gel:

- a negative control with genomic DNA prepared from a non-transgenic rice plant. This negative control is used to confirm the absence of background hybridization.
- 20 - a DNA positive control: With a heterozygous single copy integration of the transgene into the *Oryza sativa* genome, 10 µg of genomic DNA has the same number of molecule equivalents as  $\pm 19$  picogram of 1501 bp PvuI-HindIII fragment of pB5/35Sbar DNA (*Oryza sativa* diploid genome size:  $0.8 \times 10^9$  bp). The amount  
25 representing one plasmid copy per genome is added to 1 µg of digested non-transgenic *Oryza sativa* DNA. This reconstitution sample is used to show that the hybridizations are performed under conditions allowing hybridization of the probe with target sequences.

30 Phage Lambda DNA (strain Clind 1 ts 857 Sam 7, Life Technologies) digested with PstI was included as size standard.

After electrophoresis, the DNA samples (digested rice genomic DNA, controls and size standard DNA) were transferred to a Nylon membrane by capillary blotting during 12 to 16 hours. The DNA templates used for probe preparation were prepared by restriction digestion of plasmid pB5/35Sbar with EcoRI. This released a 1327 bp DNA fragment that encompasses a relevant part of the transforming DNA (1501 bp PvuI-HindIII fragment). After purification, the DNA fragment was labeled according to standard procedures, and used for hybridizing to the membrane.

Hybridization was performed under standard stringency conditions: The labeled probe was denaturated by heating for 5 to 10 minutes in a water bath at 95°C to 100°C and chilling on ice for 5 to 10 minutes and added to the hybridization solution (6 X SSC (20 X SSC is 3.0 M NaCl, 0.3 M Na citrate, pH 7.0), 5 X Denhardt's (100 X Denhardt's = 2% Ficoll, 2% Polyvinyl pyrrolidone, 2% Bovine Serum Albumin), 0.5 % SDS and 20 µg/ml denatured carrier DNA (single-stranded fish sperm DNA, with an average length of 120 - 3000 nucleotides). The hybridization was performed overnight at 65°C. The blots were washed three times for 20 to 40 minutes at 65°C, with the wash solution (2 X SSC, 0.1 % SDS).

The autoradiographs were electronically scanned.

The restriction patterns obtained after digestion of GAT-OS2 genomic DNA with different restriction enzymes is presented in Figure 1 and summarized in Table 1.

Table 1: Restriction map of GAT-OS2

Lane number	DNA loaded	Migration of hybridizing DNA fragments between size marker bands		Estimated length of the hybridizing DNA fragments.
		Larger than	Smaller than	
1	Lambda DNA with PstI	Not Applicable	Not Applicable	



2	GAT-OS2 – EcoRI	1159	1700	1327 bp (*)
3	GAT-OS2 – BamHI	1700 805	2140 1093	2,0 kbp 805 bp (*)
4	GAT-OS2 – EcoRV	5077 2838	- 4507	12 kbp 3,8 kbp
5	GAT-OS2 – HindIII	5077	11497	5,3 kbp
6	GAT-OS2 – NcoI	2838 2838	4507 4507	4,1 kbp 3,1 kbp
7	GAT-OS2 – NsiI	4749	11497	5,1 kbp
8	Non- transgenic rice	-	-	-
9	Control plasmid DNA – EcoRI	1159	1700	1327 bp

(\*) the lengths of these fragments are those predicted from the restriction map of the 1501 bp PvuI-HindIII fragment of pB5/35S.

## (2) identification of flanking sequences

5

The sequence of the regions flanking the inserted transgene in the GAT-OS2 event was determined using the thermal asymmetric interlaced (TAIL-) PCR method as described by Liu et al. (1995, The Plant Journal 8(3):457-463). This method utilizes three nested specific primers in successive reactions together with a shorter arbitrary  
 10 degenerate (AD) primer so that the relative amplification efficiencies of specific and non-specific products can be thermally controlled. The specific primers were selected for annealing to the border of the transgene and based on their annealing conditions.

A small amount (5 $\mu$ l) of unpurified secondary and tertiary PCR products were analyzed on a 1% agarose gel. The tertiary PCR product was used for preparative amplification, purified and sequenced on an automated sequencer using the DyeDeoxy Terminator cycle kit.

5

# 1. TAIL-PCR of the HindIII site

The primers used were:

10

	Sequence (5' $\rightarrow$ 3')	Position in pB5/35Sbar
Degenerate primer MDB556	CNg.ASN.AgW.TWg.CAT.A	-----
Primary TAIL MDB010	gCA.CCA.TCg.TCA.ACC.ACT.ACA.TCg	2905 $\rightarrow$ 2882
Second. TAIL MDB559	TTC.Tgg.Cag.CTg.gAC.TTC.AgC	2483 $\rightarrow$ 2463
Tertiary TAIL MDB411	Agg.CAT.gCC.gCT.gAA.ATC.ACC	2407 $\rightarrow$ 2386

whereby: N=A,C,T or g; S=C or g; W=A or T

The fragment amplified using MDB556-MDB411 was ca. 400 bp of which 113 bp were sequenced (5' flank: SEQ ID No 1). The sequence between bp 1 and bp 92 comprised plant DNA, while the sequence between bp 93 and bp 113 corresponded to pB5/35Sbar DNA.

15

# 2. TAIL-PCR of the PvuI site

The primers used were:

	Sequence (5' → 3')	Position in pB5/35Sbar
Degenerate primer MDB285	NTC.gAS.TWT.SgW.gTT	-----
Primary TAIL MDB424	AAG.gAT.AgT.ggg.ATT.gTg.Cg	3037→3056
Secondary TAIL MDB442	AAT.ggA.ATC.CgA.ggA.ggT.TTC.C	3283→3304
Tertiary TAIL MDB410	TCg.TgC.TCC.ACC.Atg.TTg.ACg	3390→3410

whereby: N=A,C,T or g; S=C or g; W=A or T

5

The fragment amplified using MDB285-MDB410 was ca. 1200 bp (3' flank: SEQ ID No 2). The sequence between bp 1 and bp 604 corresponded to pB5/35Sbar DNA, while bp 605 to bp 1279 comprised plant DNA.

10 (3) Identification of the target site deletion

Using primers corresponding to sequences within the flanking regions of the transgene on the wildtype *Oryza sativa* var. Bengal as a template, the insertion site of the transgene was identified.

15

The following primers were used:

	Sequence (5' → 3')	Position in 5'flank (SEQ ID No 1)	Position in 3'flank (SEQ ID No 2)
YTP059	TCg.gAC.AAC.CgC.gAT.AgT.TCg	56 → 76	-----
OSA04	TCg.CAT.ATg.TAT.gTA.ACA.CgC	-----	717 → 697

This yielded a 168 bp fragment (SEQ ID No 3) in which bp 38 to 55 corresponds to a target site deletion.

- 5 Thus, the complete rice insertion region (SEQ ID No 9) as sequenced comprises:
- |          |                      |                               |
|----------|----------------------|-------------------------------|
| 1-92:    | 5' flanking region   | bp 1-92 of SEQ ID No 1        |
| 93-110:  | target site deletion | bp 38 to 55 of SEQ ID No 3    |
| 111-785: | 3' flanking region   | bp 605 to 1279 of SEQ ID No 2 |

#### 10 (4) genetic analysis of the locus

The genetic stability of the insert was checked by molecular and phenotypic analysis in the progeny plants over several generations.

- 15 Southern blot analyses on glufosinate resistant plants of GAT-OS2 rice plants of the T<sub>0</sub>, T<sub>1</sub> and T<sub>2</sub> generation were compared and were found to be identical. This proves that the molecular configuration of the transgene in GAT-OS2 containing plants was stable.

- 20 The GAT-OS2 event displayed Mendelian segregation for the transgene as a single genetic locus in at least three subsequent generations indicating that the insert is stable.

On the basis of the above results GAT-OS2 was identified as an elite event.

b) Development of diagnostic tools for identity control

5

The following protocols were developed to identify any rice plant material comprising the elite event GAT-OS2.

**(1) GAT-OS2 Elite event Restriction map identification protocol**

10

Rice plants containing the elite event GAT-OS2 can be identified by Southern blotting using essentially the same procedure as described in Example 3 a) (1). Thus rice genomic DNA is 1) digested with at least three, preferably at least 4, particularly with at least 5, more particularly with all of the following restriction enzymes: EcoRI, BamHI, EcoRV, HindIII, NcoI, NsiI, 2) transferred to nylon membranes and 3) hybridized with the 1327 bp EcoRI fragment of plasmid pB5/35Sbar. If, with respect to each of the restriction enzymes used, DNA fragments are identified with the same length as those listed in Table 1, the rice plant is determined to harbor elite event GAT-OS2.

20

**(2) GAT-OS2 Elite event Polymerase Chain Reaction identification protocol**

A test run, with all appropriate controls, has to be performed before attempting to screen unknowns. The presented protocol might require optimization for components that may differ between labs (template DNA preparation, Taq DNA polymerase, quality of the primers, dNTP's, thermocycler, etc.).

25

Amplification of the endogenous sequence plays a key role in the protocol. One has to attain PCR and thermocycling conditions that amplify equimolar quantities of both the endogenous and transgenic sequence in a known transgenic genomic DNA template. Whenever the targeted endogenous fragment is not amplified or whenever

30

the targeted sequences are not amplified with the same ethidium bromide staining intensities, as judged by agarose gel electrophoresis, optimization of the PCR conditions may be required.

## 5    **Template DNA**

Template DNA is prepared according to Edwards *et al.* (Nucleic Acid Research, 19, p1349, 1991). When using DNA prepared with other methods, a test run utilizing different amounts of template should be done. Usually 50 ng of genomic template  
10    DNA yields the best results.

### **Assigned positive and negative controls**

The following positive and negative controls should be included in a PCR run:

15

- Master Mix control (DNA negative control). This is a PCR in which no DNA is added to the reaction. When the expected result, no PCR products, is observed this indicates that the PCR cocktail was not contaminated with target DNA.
- 20    - A DNA positive control (genomic DNA sample known to contain the transgenic sequences). Successful amplification of this positive control demonstrates that the PCR was run under conditions which allow for the amplification of target sequences.
- 25    - A wildtype DNA control. This is a PCR in which the template DNA provided is genomic DNA prepared from a non-transgenic plant. When the expected result, no amplification of the transgene PCR product but amplification of the endogenous PCR product, is observed this indicates that there is no detectable transgene background amplification in a genomic DNA sample.

30

## Primers

The following primers, which specifically recognize the transgene and a flanking sequence of GAT-OS2 are used:

5

OSA03: 5'-gAC.TCT.gTA.TgA.ACT.gTT.CgC-3' (SEQ ID 4)  
(target: P35S)

10

OSA04: 5'-TCg.CAT.ATg.TAT.gTA.ACA.CgC-3' (SEQ ID 5)  
(target: plant DNA)

Primers targeting an endogenous sequence are always included in the PCR cocktail. These primers serve as an internal control in unknown samples and in the DNA positive control. A positive result with the endogenous primer-pair demonstrates that there is ample DNA of adequate quality in the genomic DNA preparation for a PCR product to be generated. The endogenous primers used are:

20

OSA01: 5'-gAT.CAg.TgC.Agg.CAA.TAC.Tgg-3' (SEQ ID 6)  
(Phospholipase D gene Acc. No. AB001919, 3836→3856)

OSA02: 5'-TTC.CTA.ACA.TgT.ggg.TgT.Cg-3' (SEQ ID 7)  
(Phospholipase D gene Acc. No. AB001919, 4291→4272)

## Amplified fragments

25

The expected amplified fragments in the PCR reaction are:

For primer pair OSA01-OSA02: 457bp (endogenous control)  
For primer pair OSA03-OSA04: 313bp (GAT-OS2 Elite Event)

30

**PCR conditions**

The PCR mix for 50 $\mu$ l reactions contains:

- 5                      5  $\mu$ l template DNA  
                         5  $\mu$ l 10x Amplification Buffer (supplied with Taq polymerase)  
                         1  $\mu$ l 10 mM dNTP's  
                         1  $\mu$ l OSA01 (10pmoles/ $\mu$ l)  
                         1  $\mu$ l OSA02 (10pmoles/ $\mu$ l)  
10                     2  $\mu$ l OSA03 (10pmoles/ $\mu$ l)  
                         2  $\mu$ l OSA04 (10pmoles/ $\mu$ l)  
                         0.2  $\mu$ l Taq DNA polymerase (5 units/ $\mu$ l)  
                         water up to 50  $\mu$ l

- 15                    The thermocycling profile to be followed for optimal results is the following:

- 4 min. at 95°C  
                         Followed by: 1 min. at 95°C  
                         1 min. at 57°C  
                         2 min. at 72°C  
20                     For 5 cycles  
                         Followed by: 30 sec. at 92°C  
                         30 sec. at 57°C  
                         1 min. at 72°C  
                         For 22 to 25 cycles  
25                     Followed by: 5 minutes at 72°C

**Agarose gel analysis**

- Between 10 and 20 $\mu$ l of the PCR samples should be applied on a 1.5% agarose gel  
30 (Tris-borate buffer) with an appropriate molecular weight marker (e.g. 100bp ladder  
PHARMACIA).



### Validation of the results

5 Data from transgenic plant DNA samples within a single PCR run and a single PCR cocktail should not be acceptable unless 1) the DNA positive control shows the expected PCR products (transgenic and endogenous fragments), 2) the DNA negative control is negative for PCR amplification (no fragments) and 3) the wild-type DNA control shows the expected result (endogenous fragment amplification).

10 Lanes showing visible amounts of the transgenic and endogenous PCR products of the expected sizes, indicate that the corresponding plant from which the genomic template DNA was prepared, has inherited the GAT-OS2 elite event. Lanes not showing visible amounts of the transgenic PCR product and showing visible amounts of the endogenous PCR product, indicate that the corresponding plant from which the  
15 genomic template DNA was prepared, does not comprise the elite event. Lanes not showing visible amounts of the endogenous and transgenic PCR products, indicate that the quality and/or quantity of the genomic DNA didn't allow for a PCR product to be generated. These plants cannot be scored. The genomic DNA preparation should be repeated and a new PCR run, with the appropriate controls, has to be performed.

20

### Use of discriminating PCR protocol to identify GAT-OS2

25 Rice leaf material from plants comprising different transgenic events (samples 1 to 10) was tested according to the above-described protocol. Samples from M202 wild-type and Bengal wild-type were taken as negative controls.

The results of the PCR analysis are illustrated in Figure 2. Samples 8 and 9 (which in fact contained DNA from plants derived from the same event) are recognized as comprising elite event GAT-OS2. All other tested lines do not comprise this elite  
30 event.

Example 4: Introgression of GAT-OS2 into preferred cultivars

Elite event GAT-OS2 is introduced by repeated backcrossing into the following cultivars:

- 5 - California Temperate Japonicas (such as but not limited to M204, M202, M201, M103)
- California Tropical Japonicas (such as but not limited to L201, L202)
- Japanese and Korean Temperate Japonicas (such as but not limited to Koshihikari and Milyang)
- 10 - Australian Temperate Japonicas (such as but not limited to Millin and Jarrah)
- Mediterranean Temperate Japonicas (such as but not limited to Ballila, Arborio)
- Chinese Indicas (such as but not limited to Guichao, Congui 314, Teqing)
- Southern United State Tropical Japonicas, long grain (such as but not limited to Drew, Cypress, Jefferson, Priscilla, Cocadrie)
- 15 - Southern United State Tropical Japonicas, medium grain (such as but not limited to Bengal, Mars, Brazos, Mercury)
- South American Tropical Japonicas, long grain (such as but not limited to El Paso 144, IRGA 409)
- Far Eastern basmati and jasmine types (Kasmir, Kwao Dak Mali)
- 20 - African javanica types (bulu rices)

It is observed that the introgression of the elite event into these cultivars does not significantly influence any of the desirable phenotypic or agronomic characteristics of these cultivars (no linkage drag) while expression of the transgene, as determined by  
25 glufosinate tolerance, meets commercially acceptable levels. This confirms the status of event GAT-OS2 as an elite event.

As used in the claims below, unless otherwise clearly indicated, the term "plant" is intended to encompass plant tissues, at any stage of maturity, as well as any cells,  
30 tissues, or organs taken from or derived from any such plant, including without

limitation, any seeds, leaves, stems, flowers, roots, single cells, gametes, cell cultures, tissue cultures or protoplasts.

Seed comprising elite event GAT-OS2 was deposited as GAT-OS2 at the ATCC under  
5 number: ATCC 203352.

34  
CLAIMS

- 1) A transgenic, glufosinate tolerant rice plant, cell, tissue or seed, which is characterized in that:
- a) the genomic DNA of said plant, cell, tissue or seed is capable of yielding  
5 at least three restriction fragments or pairs of restriction fragments,  
wherein said restriction fragments or pairs of restriction fragments are  
selected from the group of:
- i) one EcoRI fragment with a length between 1159 and 1700 bp;  
ii) one pair of BamHI fragments wherein one has a length between 805  
10 and 1093 bp and the other has a length between 1700 and 2140 bp;  
iii) one pair of EcoRV fragments wherein one has a length between 2838  
and 4507 bp and the other has a length of more than 5077 bp;  
iv) one HindIII fragment with a length between 5077 and 11497 bp;  
v) one pair of NcoI fragments both with lengths between 2838 and 4507  
15 bp;  
vi) one NsiI fragment with a length between 4749 and 11497 bp;  
wherein each of said restriction fragments is capable of hybridizing, under  
standard stringency conditions, with the 1327 bp fragment obtainable by  
EcoRI digestion of the plasmid having the nucleotide sequence of SEQ ID  
20 No 8;  
and/or,
- b) a DNA fragment of between 290 and 350 bp, preferably of about 313 bp  
can be amplified from the genomic DNA of said plant, cell, tissue or seed  
using a polymerase chain reaction with two primers having the nucleotide  
25 sequence of SEQ ID No 4 and SEQ ID No 5, respectively.
- 2) The plant, cell, tissue or seed of claim 1, which is characterized in that a DNA  
fragment of between 290 and 350 bp, preferably of about 313 bp, can be  
amplified from genomic DNA of said plant, cell, tissue or seed using a  
30 polymerase chain reaction with two primers having the nucleotide sequence of  
SEQ ID No 4 and SEQ ID No 5, respectively.

- 3) The plant, cell, tissue or seed of claim 1, which is characterized in that genomic DNA of said plant, cell, tissue or seed is capable of yielding at least three restriction fragments or pairs of restriction fragments, wherein said restriction fragments or pairs of restriction fragments are selected from the group of:
- i) one EcoRI fragment with a length between 1159 and 1700 bp;
  - ii) one pair of BamHI fragments wherein one has a length between 805 and 1093 bp and the other has a length between 1700 and 2140 bp;
  - iii) one pair of EcoRV fragments wherein one has a length between 2838 and 4507 bp and the other has a length of more than 5077 bp;
  - iv) one HindIII fragment with a length between 5077 and 11497 bp;
  - v) one pair of NcoI fragments both with lengths between 2838 and 4507 bp;
  - vi) one NsiI fragment with a length between 4749 and 11497 bp;
- wherein each of said restriction fragments is capable of hybridizing, under standard stringency conditions, with the 1327 bp fragment obtainable by EcoRI digestion of the plasmid having the nucleotide sequence of SEQ ID No 8.
- 4) The plant, cell, tissue or seed of claim 3, which is characterized in that genomic DNA of said plant, cell, tissue or seed is capable of yielding at least four restriction fragments or pairs of restriction fragments selected from said group.
- 5) The plant, cell, tissue or seed of claim 4, which is characterized in that genomic DNA of said plant, cell, tissue or seed is capable of yielding at least five restriction fragments or pairs of restriction fragments selected from said group.

- 6) The plant, cell, tissue or seed of claim 5, which is characterized in that genomic DNA of said plant, cell, tissue or seed is capable of yielding the six restriction fragments or pairs of restriction fragments selected from said group.
- 5 7) The plant, cell, tissue or seed of any one of claims 3 to 6 which is further characterized in that a DNA fragment of between 290 and 350 bp, preferably of about 313 bp, can be amplified from genomic DNA of said plant, cell, tissue or seed using a polymerase chain reaction with two primers having the nucleotide sequence of SEQ ID No 4 and SEQ ID No 5, respectively.
- 10 8) A plant which is grown from a seed deposited at the ATCC under number ATCC 203352.
- 9) A cell or tissue of the plant of claim 8.
- 15 10) A seed deposited at the ATCC under number 203352
- 11) A transgenic, glufosinate tolerant rice plant of any one of claims 1 to 7 which is obtainable by propagation of, and/or breeding with, a rice plant grown from a seed deposited at the ATCC under number 203352.
- 20 12) A process for cultivating rice plants which comprises growing plants of any one of claims 1 to 8.
- 25 13) The process of claim 12 which further comprises applying a herbicide with glufosinate as an active ingredient to the cultivated rice plants.
- 14) A process for breeding rice which comprises a crossing with a plant of any one of claims 1 to 8.

- 15) A process for producing a transgenic cell of a rice plant which comprises inserting a recombinant DNA molecule into a part of the chromosomal DNA of a rice cell characterized by the sequence of SEQ ID No 9.
- 5 16) A transgenic cell of a rice plant obtainable by the method of claim 15.
- 17) A process for producing a transgenic rice plant which comprises inserting a recombinant DNA molecule into a part of the chromosomal DNA of a rice cell characterized by the sequence of SEQ ID No 9, and regeneration of a rice  
10 plant from the transformed rice cell.
- 18) A transgenic rice plant obtainable by the method of claim 17.
- 19) A method for identifying a transgenic plant, or cells or tissues thereof,  
15 comprising the elite event GAT-OS2, which method comprises establishing one or both of the following characteristics:
- a) the genomic DNA of a plant, cell, tissue or seed is capable of yielding at least three of the restriction fragments or pairs of restriction fragments selected  
20 from the group of:
- i) one EcoRI fragment with a length between 1159 and 1700 bp;
- ii) one pair of BamHI fragments wherein one has length between 805 and 1093 bp, and the other has a length between 1700 and 2140 bp;
- iii) one pair of EcoRV fragments wherein one has a length between 2838 and 4507 bp, and the other has a length of more than 5077 bp;  
25 iv) one HindIII fragments with a length between 5077 and 11497 bp;
- v) one pair of NcoI fragments both with lengths between 2838 and 4507 bp and one of about 4,1 kbp;
- vi) one NsiI fragment with a length between 4749 and 11497 bp;
- 30 wherein each of the restriction fragments is capable of hybridizing under standard stringency conditions, with the 1327 bp fragment obtainable by EcoRI digestion of the plasmid having the nucleotide sequence of SEQ ID No. 8; and/or,

- b) the genomic DNA of the plant, cell, tissue or seed can be used to amplify a DNA fragment of between 290 and 350 bp, using a polymerase chain reaction with two primers having the nucleotide sequence of SEQ ID No. 4 and SEQ ID No. 5, respectively.

5

- 20) The method of claim 19, which comprises establishing whether the genomic DNA of the transgenic plant, or its cells or tissues is capable of yielding all six of said restriction fragments or pairs of restriction fragments.

10

- 21) The method of claim 19, which comprises establishing whether the genomic DNA of the transgenic plant, or its cells or tissues can be used to amplify a DNA fragment of about 313 bp using a polymerase chain reaction with two primers having the nucleotide sequence of SEQ ID No.4 and SEQ ID No. 5, respectively.

15

- 22) A kit for identifying a transgenic plant, its cells or tissues comprising the MS-B2 elite event, said kit comprising at least two PCR probes, one of which recognizes a sequence within the foreign DNA of GAT-OS2, the other which recognizes a sequence within the 3' or 5' flanking region of GAT-OS2.

20

- 23) The kit of claim 22, said kit comprising the PCR probes having the nucleotide sequence of SEQ ID No. 4 and SEQ ID No. 5, respectively.



Figure 1





Figure 2



# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/25667

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : Please See Extra Sheet.  
US CL : 800/278, 295, 298, 320.2; 536/23.6, 24.2; 435/172.3, 419, 468  
According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 800/278, 295, 298, 320.2; 536/23.6, 24.2; 435/172.3, 419, 468

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WEST, CABA, CAPLUS, BIOSIS  
search terms: rice, gluphosinate, BAR, GAT-OS2

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,641,664 A (D'HALLUIN et al) 24 June 1997, see entire document.	1-23
Y	US 5,561,236 A (LEEMANS et al) 01 October 1996, see entire document.	1-23

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

*A*	document defining the general state of the art which is not considered to be of particular relevance	*T*	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*E*	earlier document published on or after the international filing date	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*L*	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*O*	document referring to an oral disclosure, use, exhibition or other means	*A*	document member of the same patent family
*P*	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search  
09 FEBRUARY 2000

Date of mailing of the international search report

05 APR 2000

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231  
Facsimile No. (703) 305-3230

Authorized officer

THOMAS HAAS

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)\*

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/25667

## A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

C12N 5/04, 15/00, 15/05, 15/29, 15/64, 15/82; A01H 3/00, 4/00, 5/00